

THE LINK BETWEEN DIABETES & ERECTILE DYSFUNCTION

A Patient Guide



Inside this guide

Erectile Dysfunction: It's common and treatable	6
ED and Diabetes	8
Erections and How They Work	10
What is Erectile Dysfunction?	11
Treatment Options	12
Coloplast Penile Implants	13
Frequently Asked Questions	18
Sexual Health Inventory for Men (SHIM)	20
Supporting a Partner with ED	22
Insurance Information	24
Take the Next Step	27



Let's talk about it.

You can find a solution that works.

1 in 4 men over age 65 experience some degree of erectile dysfunction (ED).¹⁶ And men with diabetes experience ED at three times the rate of non-diabetics¹ because of the damage diabetes can cause to the nerves and blood vessels that control the ability to get an erection.

Living with ED is tough. You may think that no one understands how you feel, and no one wants to talk about it. But there are millions of men who experience ED and know how it can affect daily life.

Our patient educators—real people with ED who found a solution that works for them—are ready to talk about it. They share with others their experience with ED: what it's like, what you can expect, and how they finally got back to feeling like themselves again.

Review this guide to learn more about ED and then talk with your doctor and a patient educator.

Erectile Dysfunction: It's common and treatable

If you have Erectile Dysfunction (ED), you're not alone:

Approximately **30 million men** in the U.S. suffer from ED²

50% of men with diabetes experience ED.¹⁵ ED can even be the first symptom of the disease, especially in younger men.¹⁴



1 in 4 men over age 65 experience some degree of ED¹⁷ While ED is common, it's not an inevitable part of aging, and in most cases it can be overcome.²

There are many different treatments, including pills, injections and vacuum pumps. A highly effective and satisfying solution is a penile implant.⁶ An implant involves surgery, and your doctor may recommend other options first, but a penile implant can offer a highly effective, satisfying return to an active sex life.³

There's a lot to learn, and many things to consider, so let's get started.



ED and Diabetes

Diabetes can cause nerve, blood vessel, and muscle damage that results in problems like pain, numbing or loss of sensation in the hands and feet.²⁰ These issues can also result in ED problems, because nerve signals and blood flow are necessary to the process of getting an erection. And as men with diabetes get older, ED problems become even more common.¹⁴

Finding the Right Treatment Option

Most cases of ED can be treated, and oral medications are often the first step. However, they have been shown to be less effective for men with diabetes, and more advanced treatment options may be needed.⁵ In those cases, ED treatment options such as injection therapy, vacuum devices, or a penile implant may be recommended.²

What it means for you

If you are a diabetic and experiencing ED, talk with your doctor or diabetes educator about the treatment options that may be best for you.



3x

Men with diabetes experience ED at three times the rate of other men, and it often affects them at a younger age.¹



95%

Nearly all men over age 70 are likely to have some difficulty with ED.¹⁶

35% - 75%

of men with diabetes will experience at least some degree of ED.¹⁶


Erections and How They Work²

Sexual stimulation and excitement cause the brain, nerves, heart, blood vessels and hormones to work together to produce a rapid increase in the amount of blood flowing to the penis.

The blood fills the two spongy chambers (called the corpora cavernosa) in the shaft of the penis.

As the chambers rapidly fill with blood, they expand, compressing the veins and trapping the blood in the penis. The penis becomes firm and elongated, resulting in an erection.

ED problems can happen when erectile tissue is damaged, when there isn't enough blood flow, or when your brain doesn't send the right signals – such as when nerve damage occurs from diabetes.



What is Erectile Dysfunction?2

ED is the consistent inability to sustain an erection sufficient for sexual intercourse. That can be:


- Not getting an erection at all
- Getting an erection inconsistently
- Getting an erection that doesn't stay hard enough for sex

Causes of ED

For more than 80% of men with ED, it is caused by a physical problem or disorder.⁴ The cause can usually be identified, and proper treatment can help you return to a satisfying sex life.³

Physical causes2:

- Injury (such as brain or spinal cord injury)
- Disease (such as diabetes, high blood pressure, or high cholesterol)
- Surgery (such as removal of the prostate gland)
- Substance use (such as tobacco, drugs, alcohol, and some medications)





Proper treatment can help you return to a satisfying sex life.⁶

Treatment Options²

There are different ways to treat ED. Your doctor can discuss these with you, and help you consider your individual needs, and the benefits and risks of each treatment option.

- Lifestyle changes like stopping smoking, losing weight, and eating healthier
- Oral ED medications like Viagra, Cialis or Levitra
- Vacuum erection devices
- Injecting medication directly into the penis
- Penile implant: flexible or inflatable



"I underwent all the processes I thought might help, but nothing helped me. So I said well, I have to do this because it's the only solution I have. And, the only thing I regret is not doing it sooner, because I'm a much happier man after the Coloplast Titan penile implant."

- MARTIN*


Coloplast Penile Implants

A penile implant is a discrete device that is placed into a man's penis and is designed to help him get an erection.⁷ Once implanted, it helps put the man back in control of his body, and can be used at any time, unlike other treatment options where there may be a short waiting period for them to be effective.¹⁸ A penile implant is an effective solution⁶ for men who are suffering from ED that has not been resolved by other available treatment options.

A penile implant involves a routine surgical procedure. The majority of men can return home within a day of surgery, and after full recovery, may resume sexual activity after 4-6 weeks.² Your physician will discuss the details of the procedure, your recovery process, and using the implant.

Penile implants have been an effective standard of care for men suffering from ED for many years⁶, and can offer hope and a return to intimacy and a normal life for many men with ED.¹⁹ There are two types of penile implants that Coloplast offers:

- Titan[®] inflatable penile implant produces a controllable erection that acts and feels more like a natural one.⁶
- **Genesis**° flexible penile implant produces a permanently firm penis and may be appropriate for men with limited hand dexterity.⁹



Coloplast Titan[®] Inflatable Penile Implant⁷

The Titan inflatable penile implant is a self-contained, fluid-filled system made from silicone and Bioflex*, a supple yet durable material. There are three parts to the system, connected by silicone tubing:



A reservoir (placed in the abdomen)



Two cylinders (placed in the shaft of the penis)




A pump (placed in the scrotum)

When you squeeze the pump bulb, it moves the fluid from the reservoir into the penile cylinders, creating an erection. When you press the deflate button on the pump, the fluid moves out of the penis and back into the reservoir for a natural looking flaccid state.

Visit **TitanInflate.com** for a demonstration video.

Titan Features⁶

- Easy to use and activate
- One-step deflation
- Not visibly noticeable when flaccid
- Acts and feels like a natural erection
- · Maximizes the girth of the penis



"The Coloplast Titan has brought the intimacy and spontaneity back to our love life. It was lost with the pills, pumps, and shots, which all made it too mechanical and more of a chore than an expression of our love for each other."


- RAYMOND*

Coloplast Genesis® Flexible Penile Implant8

The Genesis is a flexible penile implant that consists of two firm but flexible rods that are placed into the shaft of the penis. There are no other parts to this implant. To have an erection, you simply hold the penis and move it into the desired position. When you are finished, you bend it back down to conceal it.

Genesis Features9

- Easy for you or your partner
- · Good option for men with limited dexterity
- Because it stays firm when not erect, may be more difficult to conceal in clothing





Important Risk Information

A penile implant requires surgery. Risks of surgery may include but are not limited to complications such as infection, swelling, pain and discomfort. Once implanted, there may be a chance of infection, erosion, product migration or device malfunction which may require additional surgery. Your physician will provide complete risk information for your situation.

Frequently Asked Questions

Q. Can I have an orgasm with a penile implant?

A. You should be able to have an orgasm with a penile implant if you were able to have one before the implant.¹⁰ Consult your physician about your expected outcome.

Q. What is the recovery time?

A. Everyone is different and recovery time varies, but typically it's between 4-6 weeks until you can resume sexual activity.² Your physician will determine what you can and can't do during this time. It is important to follow all recommendations from your physician for the best outcome.

Q. Will I lose any length after getting a penile implant?

A. Each penile implant is custom-fitted to your anatomy, and the implant itself does not lead to a loss in size. Depending on your medical history, changes in anatomy or atrophy can result in loss of penile length prior to the implant, and it may not be unusual to lose 1-2 centimeters from your original erect length.¹²

Q. Will anyone notice that I have an implant?

A. The implant is completely placed inside your body. The Titan® inflatable penile implant is not visibly noticeable. The penis appears relaxed and normal when in the flaccid state6, and it is not obvious by looking at a man that he has an implant. The Genesis® flexible rod implant stays firm when not in the erect position, and it may "show" through clothes.

Q. How long does the penile implant last?

A. A study showed that 60% of initial implants would survive 15 or more years without revision or extraction.¹¹

Q. Will I be able to have spontaneous erections with a penile implant?

A. Both the Titan and the Genesis give you the ability to have an erection instantly and spontaneously, any time you want one.^{7,8} However, the implant surgery makes it impossible to ever have a "latent" or natural erection that's not dependent on the device. Therefore, you should consider carefully whether or not an implant is the right choice for you.

Q. What is the difference between the Genesis and the Titan implants?

A. Both help give you the ability to have an erection satisfactory for intercourse. The main difference is that the Genesis is a flexible (malleable) implant that consists of two firm but bendable rods that are placed into the shaft of the penis (corpora cavernosa). There are no other parts to this implant. To have an erection, you simply hold the penis and move it into the desired position. When you are finished, you return the penis to its tucked-down position. With the Titan, you inflate the cylinders by squeezing the pump bulb located in your scrotum, and deflate by pressing the release valve to return the fluid from the cylinders into the reservoir.

[&]quot;I'm able to walk into the room now and be that guy that I always wanted to be, just confident and self-assured and knowing what I want. That has made all the difference for me."

Sexual Health Inventory for Men (SHIM)¹³

Over the past 6 months:

How do you rate your confidence that you could get and keep an erection?		Very Low 1
2. When you had erections with sexual stimulation, how often were your erections hard enough for penetration (entering your partner)?	No sexual activity 0	Almost never or never
3. During sexual intercourse, how often were you able to maintain your erection after you had penetrated (entered) your partner?	Did not attempt 0	Almost never or never
4. During sexual intercourse, how difficult was it to maintain your erection to completion of intercourse?	Did not attempt 0	Extremely difficult
5. When you attempted sexual intercourse, how often was it satisfactory for you?	Did not attempt 0	Almost never or never

Low 2	Moderate 3	High 4	Very High 5
A few times (much less than half the time)	Sometimes (about half the time)	Most times (much more than half the time)	Almost always or always 5
A few times (much less than half the time)	Sometimes (about half the time)	Most times (much more than half the time)	Almost always or always 5
Very difficult 2	Difficult 3	Slightly difficult 4	Not difficult 5
A few times (much less than half the time)	Sometimes (about half the time)	Most times (much more than half the time)	Almost always or always

Add the numbers	corresponding	to	questions	1-5
TOTAL:				

The Sexual Health Inventory for Men further classifies ED severity with the following breakpoints:

1-7 Severe ED 8-11 Moderate ED 12-16 Moderate to Mild ED 17-21 Mild ED

Supporting a Partner with ED

DON'T blame yourself.

When men struggle with ED, spouses or partners tend to blame themselves first. They think it's their fault, that maybe their partner isn't attracted to them anymore. They often don't realize that the ED is likely caused by a medical problem.

DO your homework.

Many people view ED as a sexual issue, when in fact, it's usually a physical one. Conditions such as diabetes, high cholesterol, or early-stage heart conditions can all contribute to ED.² Even certain medications can bring on ED.² The sooner you realize that this is a medical condition affecting your partner's body, the faster the healing can begin.

DON'T approach the issue with negative emotions.

A man with ED can often experience deep feelings of shame, loneliness, anxiety and depression. He may often say that the inability to have an erection makes him feel like less of a man. He may be hesitant to kiss or cuddle because he is embarrassed about where it might lead. Confronting him with feelings of hurt or anger may cause him to feel attacked and withdraw even further.

"After the operation, I noticed a change in my husband in terms of his self-esteem. It's something that's made him feel comfortable and happy; he feels like a different man."

- ANABEL*

DO open the lines of communication.

Have a conversation with him – but not in the bedroom. Put some time and space between your conversation and your last sexual encounter. Make him aware of the health conditions that can cause ED, and gently suggest he see his doctor. Some men may ask you to join them at their appointment, while others may prefer to have a private conversation with their physician. Let him decide.

DON'T tell him that his ED doesn't matter.

Some people think they are being helpful by saying their partner's ED isn't a big deal. The truth is, it may matter deeply to him, and suggesting otherwise sends the message that you don't miss intimate sexual contact with him, which can be hurtful.



Insurance Information

Employer and Individual Plans

Most plans provide coverage for penile implant surgery, as long as it meets the criteria for being medically necessary.** All plans include some limitations, however, and exclusions for penile implants do occur. Before scheduling a procedure, you'll need to determine if it is covered by your plan. Your physician's office will likely contact your insurance company to check benefits, but you may wish to contact them yourself to make sure you understand what any out-of-pocket expenses might be.

Medicare

Medicare and Medicare Advantage provide coverage for penile implant surgery as long as it meets the criteria for being medically necessary.** Traditional Medicare does not require any authorization for these procedures, but if you have Medicare Advantage, your physician may need to get approval before scheduling your surgery.

Medicaid

Medicaid coverage for penile implants varies a great deal between states. Your physician should be able to advise you about local coverage, or you can contact the Medicaid program directly for answers about coverage.

Uninsured

If you do not have insurance or your existing insurance does not cover a penile implant, there are still options to consider for making the procedure more affordable. For example, you may wish to explore alternative insurance policies or financing options. Costs can vary widely between different physicians and facilities, and many health providers offer financial assistance for non-insured patients.

To learn more, talk with your doctor or visit www.coloplastmenshealth.com/insurance/ for additional insurance information.

** A penile implant is considered medically necessary when the patient's erectile dysfunction is the result of organic (rather than psychogenic) causes, and he has found non-invasive treatments (drugs, injections and/or vacuum devices) to be ineffective.



Let's Talk

Through our **Patient Educator Program**, we offer the opportunity for you to talk candidly with men who have a penile implant and want to share their experiences with others, as well as some of their partners. They're not doctors and they don't give medical advice. They are simply real people who once suffered with ED, and decided on a Coloplast penile implant to help restore their sex lives. You can set up an appointment to speak personally and confidentially with them, one-onone, about their experiences with ED and their journey to a satisfying solution.

Visit EDsolution.com and click on "Talk to a Patient" to schedule a conversation today.



EDsolution.com

Use the **physician locator** to find a physician who specializes in erectile dysfunction.

Confidentially **talk to a patient** or spouse about their experience.

View videos and animations about how a penile implant works.


Para pacientes de habla española, visite **LaBombita.com**



Take the Next Step

Discover whether a penile implant is the way for you to restore your sex life.

- Visit EDsolution.com to hear more patient stories, schedule a conversation with a patient educator, see how a penile implant works or to locate a urologist who specializes in ED.
- Take the Sexual Health Inventory for Men (SHIM) survey in this brochure, and share it with your doctor.
- Set up an appointment with your doctor to learn more about erectile dysfunction and penile implants and discuss your options.



- Malavige, S. Lasantha. Erectile dysfunction in diabetes mellitus. J Sex Med 2009; 6:1232-1247.
- Erectile dysfunction. NIH: National Institute of Diabetes and Digestive and Kidney Diseases. http://www.nlm.nih.gov/medlineplus/ erectiledysfunction.html.
- 3. Rajpurkar A, Dhabuwala CB. Comparison of Satisfaction Rates and Erectile Function in Patients Treated with Sildenafil, Intracavernous Prostaglandin E1 and Penile Implant Surgery for Erectile Dysfunction in Urology Practice. J Urol. 2003; 170:159-163.
- Khera M, Goldstein I. 2011. Erectile dysfunction. BMJ Clinical Evidence. http://www.clinicalevidence.com.
- Walsh TJ, Hotaling JM, Smith A, Saigal C, Wessells H. Men with diabetes may require more aggressive treatment for ED. Int J of Impotence Research 2013; 26:112-115
- 6. Data on file with Coloplast.
- Coloplast Titan®, Titan® OTR and Titan® Touch Inflatable Penile Prosthesis Instructions for Use. Coloplast Corp. 2016.
- Coloplast Genesis® Malleable Penile Prosthesis Instructions for Use. Coloplast Corp. 2014.
- Habous M, Malleable (Semi-Rigid) penile prosthesis. J Sex Med. 2015; 12:1984-1988.
- 10. Dealing with Erectile Dysfunction. UCLA Urology. http://urology.ucla.edu/dealing-with-erectile-dysfunction.
- 11. Wilson, Steven et al. Long-Term Survival of Inflatable Penile Prostheses: Single Surgical Group Experience with 2,384 First-Time Implants Spanning Two Decades. J. Sex Med. 2007; 4:1074-1079.
- 12. Savoie, Marc et al. A Prospective Study Measuring Penile Length in Men Treated with Radical Prostatectomy for Prostate Cancer. J Urol. 2003; 169:1462-1464.
- Medical Student Curriculum: Erectile Dysfunction. AUA. American Urological Association. http://http://www.auanet.org/education/ auauniversity/medical-student-education/medical-studentcurriculum/ed.
- Sexual and Urologic Problems of Diabetes. NIH: National Institute of Diabetes and Digestive and Kidney Diseases. http://diabetes.niddk. nih.gov/dm/pubs/sup/. Accessed May 2015.
- 15. Hatzimouratidis K, Hatzichristou D. How to treat erectile dysfunction in men with diabetes: from pathophysiology to treatment. Curr Diab Rep. 2014;14(11):545.
- 16. Erectile Dysfunction and Diabetes WebMD Medical Reference https://www.webmd.com/erectile-dysfunction/guide/ed-diabetes
- 17. Impotence. NIH: National Institutes of Health. https:consensus.nih. gov/1992/1992Impotence091html.htm.
- 18. www.viagra.com/learning/how-does-viagra-work accessed 2/1/18
- Quallich, S et al. Evaluation of Three Penile Prosthesis Pump Designs in a Blinded Survey of Practitioners. Urologic Nursing 2008; 28(2):101-108.
- Diabetic Neuropathy. https://www.mayoclinic.org/diseasesconditions/diabetic-neuropathy/symptoms-causes/syc-20371580 Accessed 2/2/18
- * Patient Educator has received compensation from Coloplast to provide their testimonial. Each person's situation is unique so your experience may not be the same.



Ostomy Care Continence Care Wound & Skin Care Urology Care

Coloplast Corp. Minneapolis, MN 55411 / Urology Care Surgical Support 800.258.3476 www.coloplast.us The Coloplast logo is a registered trademark of Coloplast A/S. © 2018. All rights reserved.